arXiv:1209.2450v1 [math.DG] 11 Sep 2012

SEIFERT FIBERINGS AND COLLAPSING OF INFRASOLV SPACES

September 13, 2012

OLIVER BAUES AND WILDERICH TUSCHMANN

Abstract. We give a purely geometrical smooth characterization of closed infrasolv manifolds and orbifolds by showing that, up to diffeomorphism, these are precisely the spaces which admit a collapse with bounded curvature and diameter to compact flat orbifolds. Moreover, we distinguish irreducible smooth fake tori geometrically from standard ones by proving that the former have non-vanishing D-minimal volume.

1. Introduction

In this article we consider smooth orbifolds $\mathcal{O} = X/\Gamma$, where X is a simply connected Riemannian manifold and Γ is a properly discontinuously acting group of isometries. We also assume that a finite orbifold covering space of $\mathcal{O}$ is a manifold. If X is contractible, then $\mathcal{O}$ is called an *aspherical orbifold*. For the orbifolds considered here, the notion of smooth map and diffeomorphism can be declared by considering equivariant smooth maps on X. In what follows we are interested in geometrical properties of orbifolds which distinguish their diffeomorphism class.

Our first result relates collapsing of Riemannian orbifolds and the theory of Seifert fiberings:

Theorem 1.1. *Let M be a closed Riemannian orbifold which admits a bounded curvature collapse to a compact aspherical Riemannian orbifold $\mathcal{O}$. Then M admits the structure of a smooth Seifert fibering over $\mathcal{O}$ with infranil fiber. In particular, M is an aspherical Riemannian orbifold.*

Seifert fiber spaces with infranil fiber are a special class of orbibundles whose fibers are naturally diffeomorphic to infranil-manifolds. The notion of Seifert fiber space with infranil fiber was systematically developed by Raymond, Lee and others. (See [17, 18] for recent accounts.

The precise definition of Seifert fiber spaces is also given in Section 2.3 below.)

A Riemannian orbifold $\mathcal{S} = X/\Gamma$ is called an *infrasolv orbifold* if X is isometric to a simply connected solvable Lie group S with left-invariant metric and Γ is contained in the group of affine isometries of S. It is called *infranil* if S is nilpotent. Remarkably, as is proved in [1], infrasolv orbifolds are diffeomorphic if and only if their fundamental groups are isomorphic.

Theorem 1.2. *Let $\mathcal{S}$ be a closed infrasolv manifold (or orbifold) with fundamental group $\pi_1(\mathcal{S})$. Then $\mathcal{S}$ admits a canonical Seifert fibering with infranil fiber over a compact flat orbifold $\mathcal{O}$.*

Contracting the fibers of the above bundle to a point in an inhomogeneous way (see, for example, [22]) shows that $\mathcal{S}$ admits a bounded curvature collapse to the flat orbifold $\mathcal{O}$. Moreover, in this setting $\mathcal{S}$ is infranil if and only if it collapses to a point.

We also prove the following converse to Theorem 1.2:

Theorem 1.3. *Assume that the closed manifold (or compact orbifold) M allows a bounded curvature collapse to a compact infrasolv orbifold $\mathcal{O}$. Then M is diffeomorphic to an infrasolv manifold (respectively, orbifold) $\mathcal{S}$.*

We thus obtain a purely geometrical smooth characterization of infrasolv manifolds as follows:

Corollary 1.4. *A closed manifold is diffeomorphic to an infrasolv manifold if and only if it admits a bounded curvature collapse to a compact flat orbifold.*

The corollary generalizes the famous almost flat manifold theorem of Gromov-Ruh [14, 21] from infranil to infrasolv manifolds. It also holds for orbi- instead of manifolds (cf. Section 4), and strengthens the *topological* characterizations of infranil and infrasolv manifolds in terms of collapsing, which were developed in [6] and in [22], to the smooth case.

The proof of the preceding results relies on techniques in the theory of collapsing of Riemannian manifolds (see the appendix of this article) and, as another crucial ingredient, on the rigidity of smooth Seifert fiberings with infranil fiber as established in [16]. We recall the required rigidity results in Section 2.4.

The following applications shed some light on the geometry of fake tori. Recall that a smooth fake torus $\mathcal{T}^n$ is a smooth manifold which is homeomorphic but not diffeomorphic to a standard torus T^n. The existence of smooth fake tori, $n \geq 5$, was established by Wall and Browder, see [3], [23, 15 A]. Since, as follows from [1], smooth fake tori do not carry an infrasolv structure, we first have:

Corollary 1.5. *A smooth fake torus does not allow a bounded curvature collapse to a compact flat orbifold.*

Gromov has defined the *minimal volume* of a smooth manifold M, $\mathrm{MinVol}(M)$, as the infimum of all volumes $vol_g(M)$, where g ranges over all smooth complete Riemannnian metrics on M whose sectional curvature is bounded in absolute value by one. Gromov's critical volume conjecture asserts that there exists $\delta(n) > 0$ such that if a closed smooth n-manifold M^n admits a metric with $|\sec(M, g)| \leq 1$ and $\mathrm{vol}(M, g) \leq \delta(n)$, then $\mathrm{MinVol}(M) = 0$. For $n \leq 4$ this conjecture is known to hold, see [20, 5] and the further references cited there.

Given a real number $D > 0$, one may also consider the *D-minimal volume*, D-$\mathrm{MinVol}(M)$, where one requires that the infimum is taken over all metrics g as above, which additionally have diameter bounded from above by D. Cheeger and Rong have shown that there exists $\delta = \delta(n, D) > 0$ such that if a closed smooth n-manifold M admits a Riemannian metric g with $|\sec(M, g)| \leq 1$, $\mathrm{diam}(M, g) \leq D$, and $\mathrm{vol}(M, g) \leq \delta(n, D)$, then D-$\mathrm{MinVol}(M) = 0$, see [5]. In the special context of aspherical M this was proved by Fukaya in [7].

Call a fake smooth torus *irreducible* if it is not a product of a standard torus and a fake torus of lower dimension. For example, smooth fake tori obtained by taking the connected sum of standard tori with exotic spheres are always irreducible. Then the following holds:

Corollary 1.6. *Let $\mathcal{T}$ be a smooth fake torus which is irreducible. Then, for all $D > 0$, there exists a positive constant $\nu(D)$ such that*

$$D\text{-}\mathrm{MinVol}(\mathcal{T}) \geq \nu(D) > 0 .$$

Notice that closed infrasolvmanifolds have vanishing minimal volume, since, by Corollary 1.4, they actually have vanishing D-minimal volume for appropiate $D > 0$.

Question 1.7. *Do smooth fake irreducible tori also always have non-vanishing minimal volume ?*

2. Seifert fiber spaces with nil-geometry

2.1. **Infra-G manifolds.** Let G be a Lie group. We let $\mathrm{Aff}(G) = G \cdot \mathrm{Aut}(G)$ denote the group of affine transformations of G. *Note that the affine group* $\mathrm{Aff}(G)$ *is precisely the normalizer of (the left-action of)* G *in the group of all diffeomorphisms* $\mathrm{Diff}(G)$.

Definition 2.1. Let $\Delta \leq \mathrm{Aff}(G)$ be a discrete subgroup whose homomorphic image in $\mathrm{Aut}(G)$ has compact closure. The quotient G/Δ is called an *infra-G space*. If G/Δ is a manifold then it is called an *infra-G manifold*.

Example (Infranil manifolds)**.** Let N be a simply connected nilpotent Lie group. Then a compact infra-N manifold N/Δ is called an *infranil manifold*. For an infranil manifold N/Δ, the intersection $\Delta_0 = N \cap \Delta$ is the maximal nilpotent normal subgroup of Δ and it has finite index in Δ. (Bieberbach's first theorem, see [19, Chapter VIII].) In particular, for every infranil manifold, Δ has finite image in $\mathrm{Aut}(N)$, and N/Δ has a canonical finite normal covering by the *nilmanifold* N/Δ_0.

2.1.1. *Affine group.* Let ∇^{can} denote the natural flat connection on G which is defined by the property that left-invariant vector fields are parallel. The group of connection preserving diffeomorphisms $\mathrm{Aff}(G, \nabla^{\mathrm{can}})$ then coincides with the group of affine transformations $\mathrm{Aff}(G)$ of G. In particular, every infra-G manifold

$$G/\Delta$$

carries a flat connection ∇^{can} induced from G. The group of connection preserving diffeomorphims of $(G/\Delta, \nabla^{\mathrm{can}})$ arises from the normalizer $N_{\mathrm{Aff}(G)}(\Delta)$ of Δ in the group $\mathrm{Aff}(G)$ as

$$\mathrm{Aff}(G/\Delta, \nabla^{\mathrm{can}}) \cong N_{\mathrm{Aff}(G)}(\Delta)/\Delta\,.$$

(We will also use the shorthand $\mathrm{Aff}(G/\Delta) = \mathrm{Aff}(G/\Delta, \nabla^{\mathrm{can}})$.)

2.2. **Infra-N bundles.** Let $f : M \to B$ be a (locally trivial) fibration of smooth manifolds with fiber F.

Definition 2.2. The fibration f is called an *infra-G bundle* over B if $F = G/\Delta$ is an infra-G manifold and the structure group of f is contained in the affine group $\mathrm{Aff}(G/\Delta)$.

2.2.1. *Fiberwise affine maps.* Let $f : M \to B$ and $f' : M' \to B'$ be infra-G bundles with fiber G/Δ. Then a map

$$\varphi : M \longrightarrow M'$$

is called *fiberwise affine* if it is a morphism of $\mathrm{Aff}(G/\Delta)$-bundles (that is, equivalently φ is a bundle map which induces an affine diffeomorphism on each fiber). We let $\mathrm{Aff}(M, G/\Delta) = \mathrm{Aff}(M, G/\Delta, f)$ denote the group of all fiberwise affine diffeomorphisms of f.

2.3. **Seifert fiber spaces.**

The following notion of Seifert fiber space has been developed in [16], see also [17].

2.3.1. *Construction.* Let N be a Lie group which acts properly and freely on the manifold X, and put $W = X/N$. We assume that W is simply connected. The normalizer of N in $\mathrm{Diff}(X)$ will be denoted by $\mathrm{Diff}(X, N)$. Let π be a group and

$$\rho : \pi \longrightarrow \mathrm{Diff}(X, N)$$

an action of π which is properly discontinuous. Put $\pi_N = \rho^{-1}(N)$ for the subgroup of all elements in π which act on X via translations of $N \leq \mathrm{Diff}(X)$.

Definition 2.3. Data (X, N, π) as above define a *Seifert fiber space* if the following conditions are satisfied:

1. $\Gamma = \rho(\pi) \cap N$ is a discrete uniform subgroup of N,
2. the induced action of $\Theta = \pi\big/\pi_N$ on W is properly discontinuous.

In the situation of Definition 2.3, ρ is called a *Seifert action.* If the action of π is faithful then it gives rise to an exact sequence of groups

$$1 \longrightarrow \Gamma \longrightarrow \pi \longrightarrow \Theta \longrightarrow 1 \ \ ,$$

where $\Gamma = \pi_N$ is isomorphic to a lattice in N. *Note that, in general, the induced action of* Θ *on* W *may have a finite kernel.*

2.3.2. *Seifert bundle maps.* A Seifert fiber space gives rise to a Seifert bundle map

$$\sigma : X\big/\pi \longrightarrow W\big/\pi = W\big/\Theta$$

whose fibers are compact infra-N spaces. The space $X\big/\pi$ is called a *Seifert bundle* over the base orbifold $W\big/\Theta$ with typical fiber the N-manifold $N/\rho(\pi) \cap N$.

Remark. If X/π is a manifold then, in fact, all fibers of σ are (compact) infra-N manifolds. If also the base W/π is a manifold then all fibers of σ are naturally diffeomorphic to N/Δ, where $\Delta \leq \pi$ is the normal subgroup of π which acts trivially on W. In this situation, since π normalizes N and Δ, the Seifert bundle map σ defines a fibration whose structure group is discrete and contained in $\mathrm{Aff}(N/\Delta)$. In particular, σ is an infra-N bundle with fiber N/Δ in the sense of Definition 2.2. In general, a Seifert fibering does *not* give a locally trivial fiber bundle over the orbifold W/π, but induces the structure of an *orbibundle* over W/π. See [17] for various examples.

2.3.3. *Seifert structures on manifolds and orbifolds.*

Definition 2.4. Let M be a smooth orbifold and (X, N, π) a Seifert fiber space. A *Seifert fiber structure* on M is a diffeomorphism

$$\varphi : X/\pi \longrightarrow M\,.$$

2.4. **Seifert fiber spaces with nil-geometry.** In this subsection let N denote a simply connected nilpotent Lie group. A *lattice* in N is a discrete uniform subgroup of N.

2.4.1. *Existence.* Let Γ be a finitely generated torsion-free nilpotent group and

$$1 \longrightarrow \Gamma \longrightarrow \pi \longrightarrow \Theta \longrightarrow 1 \tag{1}$$

an exact sequence of groups.

Definition 2.5. A Seifert fiber space (X, N, π) is said to realize the group extension (1) if $\rho(\Gamma)$ is a lattice of N.

Remark. In this situation, Γ is a subgroup of finite index in $\pi_N = \rho^{-1}(N)$, and therefore the induced action of Θ on $W = X/N$ is properly discontinuous.

Theorem 2.6 ([16, §2.2])**.** *Assume that Θ acts properly discontinuously on a simply connected manifold W and let N be a simply connected nilpotent Lie group which contains Γ as a lattice. Then there exists a Seifert fiber space (X, N, π), which realizes the group extension (1) and which induces the given action of Θ on W.*

2.4.2. *Rigidity.* Let N and N' be simply connected nilpotent Lie groups, (X, N, π) and (X', N', π') Seifert fiber spaces, and $\phi : \pi \longrightarrow \pi'$ a homomorphism of groups.

Definition 2.7. An equivariant diffeomorphism of actions

$$(f, \phi) : (X, \pi) \longrightarrow (X', \pi')$$

is called an *affine equivalence* of Seifert actions if $fNf^{-1} = N'$.

Now consider an isomorphism of group extensions of the form

$$\begin{tikzcd}
1 \arrow[r] \arrow[d] & \Gamma \arrow[r] \arrow[d, "\phi_1"] & \pi \arrow[r] \arrow[d, "\phi"] & \Theta \arrow[r] \arrow[d, "\bar{\phi}"] & 1 \arrow[d] \\
1 \arrow[r] & \Gamma' \arrow[r] & \pi' \arrow[r] & \Theta' \arrow[r] & 1
\end{tikzcd} \tag{2}$$

The Seifert rigidity for nil-geometry may be stated as follows:

Theorem 2.8 ([16, §2.4])**.** *Suppose* (X, N, π) *and* (X', N', π') *are Seifert fiber spaces which realize the group extensions in diagram* (2)*. If*

$$(\bar{f}, \bar{\phi}) : (W, \Theta) \to (W', \Theta') ,$$

is an equivariant diffeomorphism then there exists a lift of $\bar{f}$ *to an affine equivalence of Seifert actions*

$$(f, \phi) : (X, \pi) \to (X', \pi') .$$

Note that, as a consequence, the Seifert rigidity constructs a diffeomorphism of Seifert fiber spaces

$$X/\pi \to X'/\pi' ,$$

which restricts to affine (in particular, connection preserving) maps on the infranil fibers.

2.5. Infranil bundles over aspherical manifolds.

Let N be a simply connected nilpotent Lie group. In this subsection we relate the notions of infra-N bundle and Seifert fiber space more closely. Recall that a manifold is called *aspherical* if its universal covering manifold is contractible. The following result asserts that over an aspherical base the structure group of an infra-N bundle may be reduced to a discrete group. This implies that *infranil bundles over an aspherical base are fiberwise affinely equivalent to a Seifert fiber space.*

Proposition 2.9. *Let* $f : M \to B$ *be an infra-*N *bundle, where* B *is an aspherical manifold and* N *is a simply connected nilpotent Lie group. Let* $\pi = \pi_1(M)$ *and assume also that* f *has compact fiber* N/Δ*. Then there exists a Seifert fiber space* (X, N, π) *which admits a fiberwise affine diffeomorphism*

$$\varphi : X/\pi \longrightarrow M$$

to the infra-N *bundle* M*.*

Proof. By the long exact homotopy sequence of the fibration f we have an exact sequence

$$1 \to \Delta \to \pi_1(M) \to \Theta \to 1$$

where $\Theta \cong \pi_1(B)$. Let $\mathsf{p} : W \to B$ be the universal covering of B, and $\bar{f} : M' \to W$ the induced infra-N bundle. The induced morphism of bundles $M' \to M$ is a covering; its group of deck transformations is isomorphic to Θ and it acts by fiberwise affine maps on $\bar{f} : M' \to W$.

Since W is contractible, the covering homotopy theorem implies that $\bar{f} : M' \to W$ is G-equivalent to the product bundle $N/\Delta \times W$, where $G = \mathrm{Aff}(N/\Delta)$ is the structure group of $\bar{f}$. That is, there exists a fiberwise affine diffeomorphism

$$N/\Delta \times W \to M'$$

of bundles over W. This map lifts to a diffeomorphism

$$\tilde{\varphi} : X \to \tilde{M}$$

of $X = N \times W$ to the universal cover $\tilde{M}$ of M. Via $\tilde{\varphi}$, the group $\pi = \pi_1(M)$ has an induced fiberwise affine action on the (trivial) N-principal bundle $X = N \times W$. Let Δ_0 be the maximal nilpotent normal subgroup of Δ. Then Δ_0 is normal in π, and also $\Delta_0 = \Delta \cap N$ is a lattice in N. Since π normalizes the lattice Δ_0, the action of π on X being fiberwise affine, Lemma 2.10 implies that π normalizes N, that is, $\pi \leq \mathrm{Diff}(N \times W, N)$. Therefore, the data $(N \times W, N, \pi)$ define a Seifert fiber space which is affinely equivalent to the infra-N bundle M. ☐

We remark:

Lemma 2.10. *Let (X, N) be a principal bundle and $\Delta_0 \leq N$ a lattice in N. Let $N_{\mathrm{Diff}(X)}(\Delta_0)$ denote the normalizer of the left action of Δ_0 in* $\mathrm{Diff}(X)$. *Then*

$$\mathrm{Aff}(X, N) \cap N_{\mathrm{Diff}(X)}(\Delta_0) \ \leq \ \mathrm{Diff}(X, N) \ .$$

Proof. Using local trivializations we may assume that $X = N \times W$ is a trivial bundle. Every element γ of $\mathrm{Aff}(X, N)$, being a fiberwise affine diffeomorphism of (X, N), gives rise to a family of automorphisms $\ell_w \in \mathrm{Aut}(N)$, $w \in W$, which is defined by the property that

$$\gamma n \gamma^{-1}(v, w) = (\ell_w(n)\, v, \bar{\gamma} w) \quad \text{, for all } n, v \in N.$$

(Here $\bar{\gamma} : W \to W$ denotes the map which γ induces on W.) If γ normalizes Δ_0 then, by Malcev rigidity of the lattice Δ_0, $\ell_w = \ell$ is constant, where $\ell \in \mathrm{Aut}(N)$ is the unique extension of the automorphism which γ induces on Δ_0. Therefore γ normalizes N. ☐

In our context, Seifert fiber spaces may be viewed as infra-N bundles with singular fibers:

Proposition 2.11. *Let $f : M_0 \to B$ be an infra-N bundle over the aspherical base manifold B, where N is a simply connected nilpotent Lie group. Assume that f has compact fiber N/Δ and let $\mu \leq \mathrm{Aff}(M_0, N/\Delta)$ be a finite group of fiberwise affine diffeomorphisms. Put*

$$M = M_0/\mu \ .$$

Let $\pi = \pi_1(M)$ be the orbifold fundamental group of M and $\pi_0 = \pi_1(M_0)$. Then there exists a Seifert fiber space (X, N, π) over the base orbifold B/μ, a Seifert fiber structure $\varphi : X/\pi \longrightarrow M$ and a commutative diagram

$$\begin{tikzcd}
X/\pi_0 \arrow[r, "\varphi_0"'] \arrow[d] & M_0 \arrow[r] \arrow[d] & B \arrow[d] \\
X/\pi \arrow[r, "\varphi"'] & M \arrow[r] & B/\mu
\end{tikzcd} ,$$

where φ_0 is a fiberwise affine diffeomorphism of bundles over B.

Proof. Let $\pi = \pi_1(M)$. By construction, the group π acts as a group of decktransformations of the universal covering $\tilde{M} \to M$ on $\tilde{M}$. This extends the corresponding action of π_0, and also induces the action of μ on $M_0 = \tilde{M}/\pi_0$. Since the fibration f is μ-equivariant, the decktransformation group $\Delta \leq \pi_0$ for the fiber of f is normalized by π, and also the maximal nilpotent normal subgroup Δ_0 is normalized by π. Recall that Δ_0 is a lattice in N.

Proposition 2.9 constructs a Seifert fiber space (X, N, π_0) and a fiberwise affine diffeomorphism φ_0 as required. Let $\tilde{\varphi} : (X, N) \to \tilde{M}$ be a lift of the diffeomorphism φ_0 to the universal covers. Then $\tilde{\varphi}$ is an affine map of bundles. We may pull back the action of π to an action on X. Since by iii) above, μ acts on the fibers of f by affine transformations, the transported action of π on X is fiberwise affine for the bundle (X, N). That is, π is contained in $\mathrm{Aff}(X, N)$. Using Lemma 2.10, it follows that, a fortiori, π is contained in $\mathrm{Diff}(X, N)$. Therefore, the data (X, N, π) satisfy the axioms of a Seifert action. □

3. Collapsing and Seifert fiberings

We consider a compact aspherical orbifold $\mathcal{O}$, where

$$\mathcal{O} = Y/\Theta \ ,$$

Y is a contractible Riemannian manifold and $\Theta \leq \mathrm{Isom}(Y)$ acts properly discontinuously on Y. Assume further that (M, g) is a closed

Riemannian orbifold which *collapses* to $\mathcal{O}$ with bounded curvatures. This means that there exists a sequence of Riemannian metrics g_i on M with uniformly bounded sectional curvature such that the metric spaces $M_i = (M, g_i)$ converge to $\mathcal{O}$ in the Gromov-Hausdorff distance. That is, we have

$$\lim_{i\to\infty} d_{\mathrm{G}H}(M_i, \mathcal{O}) = 0 \; .$$

For the Gromov-Hausdorff distance and its generalizations see Appendix A.

In this section, we show that the orbifold M carries a Seifert fiber structure over the base $\mathcal{O}$. This proves Theorem 1.1 in the introduction. In Section 3.3 we also prove Corollary 1.6.

3.1. **Equivariant collapsing on finite covers.**

Let Θ_0 be a torsion-free normal subgroup of Θ, which is of finite index. By our general assumption on orbifolds, such a subgroup of Θ always exists. Then

$$T = Y/\Theta_0$$

is a closed Riemannian manifold and it is an orbifold covering space of $\mathcal{O}$. Note that $\mathcal{O}$ is a quotient of T by the group of decktransformations of the map $T \to \mathcal{O}$. This is a finite group isomorphic to Θ/Θ_0.

Proposition 3.1. *With respect to an appropiate choice of* Θ_0*, there exists a sequence of Riemannian manifolds* $\bar{M}_i = (\bar{M}, g_i)$*, Riemannian orbifold covering maps*

$$\bar{M}_i = (\bar{M}, g_i) \longrightarrow (M, g_i),$$

and a finite group μ *which acts by isometries on* $\bar{M}_i$ *and* $T = Y/\Theta_0$ *such that*

1. $M_i = \bar{M}_i/\mu$ *and* $\mathcal{O} = T/\mu$*;*
2. *the equivariant Hausdorff distance with respect to* μ *satisfies*

$$\lim_{i\to\infty} d_{\mu-\mathrm{GH}}(\bar{M}_i, T) = 0 \; .$$

Proof. Let X_i denote the universal cover of M_i. Then $M_i = X_i/\pi$, where π acts on X_i as a discrete group of isometries. By Proposition A.7 in Appendix A, there exists a sequence $(X_i, \pi, p_i) \in \mathcal{M}^{Groups}$ such that

$$\lim_{i\to\infty} (X_i, \pi, p_i) = (Y, \Theta, q)$$

with respect to the equivariant Gromov-Hausdorff convergence with base point. Note that this implies (compare Definitions A.2, A.3.)

$$\lim_{i\to\infty} (X_i, p_i) = (Y, q) \; .$$

Now let π' be a torsion-free finite index normal subgroup of π. Using Proposition A.4, we infer that there exists a closed subgroup $\Theta' \leq \mathrm{Isom}(Y)$ such that (up to taking a subsequence)

$$\lim_{i\to\infty} (X_i, \pi', p_i) = (Y, \Theta', q) .$$

In particular, by Proposition A.6 and taking into account that the involved spaces are compact, we have

$$\lim_{i\to\infty} X_i/\pi' = Y/\Theta' .$$

Let $\mu' = \pi/\pi'$. Then μ' acts isometrically as a finite group of deck-transformations of the maps $X_i/\pi' \to X_i/\pi$ and furthermore

$$X_i/\pi \;=\; (X_i/\pi')/\mu' .$$

Let $\mathcal{M}(\mu')$ denote the space of isometry classes of compact metric spaces with isometric action of μ'. Since μ' is finite, the equivariant version of the pre-compactness theorem holds with respect to $\mathcal{M}(\mu')$ (compare Proposition A.5 in Appendix A). Therefore, after taking a subsequence, we may assume that there exists

$$\lim_{i\to\infty} (X_i/\pi', \mu') \;=\; (\mathcal{O}', \mu') \;\in \mathcal{M}(\mu').$$

This of course implies that

$$Y/\Theta = \lim_{i\to\infty} X_i/\pi = \lim_{i\to\infty} (X_i/\pi')/\mu' \;=\; \mathcal{O}'/\mu' .$$

We deduce, in particular, that $\mathcal{O}' = Y/\Theta'$ is a finite orbifold covering space of Y/Θ and Θ' is a normal subgroup of Θ. We have thus constructed a sequence of finite covering manifolds $M_i' = X_i/\pi'$ of M_i, which converges to an orbifold covering space $\mathcal{O}' = Y/\Theta'$ of $\mathcal{O}$ in the Gromov-Hausdorff-distance.

Observe that the action of π' on X_i is free, since π' is torsion-free. Therefore, we may apply Proposition A.8 in Appendix A to a torsion-free finite index normal subgroup Θ_0 of Θ', and infer that there exists a subgroup π_0 of π' such that

(1) π_0 is normal and of finite index in π', and furthermore
(2) $\lim_{i\to\infty}(X_i, \pi_0, p_i) = (Y, \Theta_0, q)$.

This gives rise to a sequence of Riemannian manifolds

$$\bar{M}_i = X_i/\pi_0$$

converging to the manifold $T = Y/\Theta_0$ in the Gromov-Hausdorff distance, that is,

$$\lim_{i\to\infty} \bar{M}_i \;=\; T .$$

Note that, by passing to a finite index subgroup of π_0, and arguing as in the first part of the proof, we can furthermore assume that

(3) π_0 is normal in π.

Now let $\mu = \pi/\pi_0$. Then μ acts on $\bar{M}_i$ by isometries and $M_i = \bar{M}_i/\mu$. Furthermore, there exists a sequence $(\bar{M}_i, \mu) \in \mathcal{M}(\mu)$ such that

$$\lim_{i\to\infty} (\bar{M}_i, \mu) = (T, \mu) \ .$$

In other words, $\lim_{i\to\infty} d_{\mu-\mathrm{GH}}(\bar{M}_i, T) = 0$. This also implies ordinary Gromov-Hausdorff convergence of $\bar{M}_i/\mu$ to T/μ. Since

$$\lim_{i\to\infty} \bar{M}_i/\mu = \lim_{i\to\infty} M_i = \mathcal{O} \ ,$$

we conclude that $\mathcal{O} = T/\mu$. This proves the proposition. □

Proposition 3.1 allows to apply equivariant collapsing results for Riemannian manifolds. Indeed, Theorem B.2 in Appendix B implies that there exist a sequence of μ-equivariant fibrations

$$f_i : \bar{M}_i \to T$$

such that, for i sufficiently large,

i) each fiber $f_i^{-1}(p)$ carries a flat connection (depending smoothly on $p \in T$), such that $f_i^{-1}(p)$ is affinely diffeomorphic to an infranil manifold N_i/Δ_i with canonical flat connection ∇^{can};

ii) the structure group of f_i is contained in the Lie group

$$\frac{\mathbb{C}(N_i)}{\mathbb{C}(N_i) \cap \Delta_i} \rtimes \mathrm{Aut}(\Delta_i) \, ,$$

where $\mathbb{C}(N_i)$ denotes the center of N_i;

iii) The affine structures on the fibers of f_i are μ-invariant;

iv) f_i is an almost Riemannian submersion.

Note, in particular, that in the terminology of Section 2.3, by ii), f_i is an infra-N_i bundle over the base T with fiber the infranil manifold N_i/Δ_i, and, by iii), μ acts by fiberwise affine maps on the bundle f_i.

3.2. **Induced Seifert fiber space structure.**

Since T is aspherical, on the level of fundamental groups the long exact homotopy sequence for the fibration $f_i : \bar{M}_i \to T$ gives a short exact sequence

$$1 \longrightarrow \Delta_i \longrightarrow \pi_0 \longrightarrow \Theta_0 \longrightarrow 1 \ \ , \tag{3}$$

where $\Delta_i \cong \pi_1(N_i/\Delta_i)$ is (isomorphic to) a subgroup of $\mathrm{Aff}(N_i)$. Since the fiber N_i/Δ_i of f_i is infranil, Δ_i intersects N_i in a uniform discrete subgroup Δ_{i0}, which is of finite index in Δ_i. In particular, Δ_i is a torsion-free finitely generated virtually nilpotent group, and Δ_{i0} is its maximal nilpotent normal subgroup. Since Δ_{i0} is normal in π_0, we obtain another short exact sequence

$$1 \longrightarrow \Delta_{i0} \longrightarrow \pi_0 \longrightarrow \bar{\Theta}_0 \longrightarrow 1 \quad , \tag{4}$$

where $\bar{\Theta}_0$ acts properly discontinuously on Y with quotient T. Proposition 2.9 and its proof now imply:

Proposition 3.2. *The infra-N_i bundle $f_i : \bar{M}_i \to T$ together with its flat fiber connections arises from a Seifert fiber construction* (X_i, N_i, π_0) *for the exact sequence* (4) *over the base T.*

This means that, for all i, there exists a Seifert fiber space (X_i, N_i, π_0) over the base $T = Y/\bar{\Theta}_0$, which realizes the group extension (4), and which has as (so called) typical fiber the nilmanifold N_i/Δ_{0i}. Moreover, there exists a diffeomorphism

$$\bar{\varphi}_i : X_i/\pi_0 \longrightarrow \bar{M}_i$$

which induces a commutative diagram

$$\begin{tikzcd}
N_i/\Delta_i \arrow[r, "\approx"] \arrow[d] & f_i^{-1}(p) \arrow[d] & \\
X_i/\pi_0 \arrow[r, "\approx", "\bar{\varphi}_i"'] \arrow[d] & \bar{M}_i \arrow[r] \arrow[d, "f_i"] & \bar{M}_i/\mu = M_i \arrow[d] \\
Y/\pi_0 \arrow[r, "\approx"] & T \arrow[r] & T/\mu = \mathcal{O}
\end{tikzcd} \quad . \tag{5}$$

such that the fiber maps $N_i/\Delta_i \overset{\approx}{\to} f_i^{-1}(p)$ are affine diffeomorphisms. Since f_i is μ-equivariant, it descends to the projection map $M_i \to \mathcal{O}$.

Let $\tilde{\varphi}_i : X_i \to \tilde{M}_i$ be a lift of $\bar{\varphi}_i$ to the universal covers. Since by iii) above, μ acts on the fibers of f_i by affine transformations, the transported action of π on X_i is fiberwise affine for the bundle (X_i, N_i). That is, π is contained in $\mathrm{Aff}(X_i, N_i)$. In the view of Proposition 2.11 we thus have:

Proposition 3.3. *The Seifert action of π_0 on (X_i, N_i) extends to a Seifert action of π on (X_i, N_i), such that the diffeomorphism $\tilde{\varphi}_i$ descends to a diffeomorphism of orbifolds $\varphi_i : X_i/\pi \to M_i$.*

Since Δ_i is a normal subgroup of π and Δ_{i0} is characteristic in Δ_i, we have induced exact sequences of the form

$$1 \longrightarrow \Delta_{i0} \longrightarrow \pi \longrightarrow \bar{\Theta}_i \longrightarrow 1 \ ; \tag{6}$$

moreover, for each i, there is a commutative diagram:

$$\begin{tikzcd}
1 \arrow[r] \arrow[d] & \Delta_{i0} \arrow[r] \arrow[d, "\shortparallel"] & \pi_0 \arrow[r] \arrow[d] & \bar{\Theta}_0 \arrow[r] \arrow[d] & 1 \arrow[d] \\
1 \arrow[r] \arrow[d] & \Delta_{i0} \arrow[r] \arrow[d] & \pi \arrow[r] \arrow[d, "\shortparallel"] & \bar{\Theta}_i \arrow[r] \arrow[d] & 1 \arrow[d] \\
1 \arrow[r] & \Delta \arrow[r] & \pi \arrow[r] & \Theta \arrow[r] & 1
\end{tikzcd} \quad . \tag{7}$$

Note that the quotient $\bar{\Theta}_i$ maps surjectively onto Θ. Since, Δ_{i0} is of finite index in Δ, $\bar{\Theta}_i$ acts properly discontinuously on $Y = X/N_i$. As diagram (5) shows, the quotient space

$$Y/\bar{\Theta}_i = Y/\Theta$$

is diffeomorphic to the orbifold $\mathcal{O} = Y/\Theta$. Therefore, the Seifert action (X_i, N_i, π) realizes the group extension (6) over the orbifold $\mathcal{O}$. We arrive at:

Theorem 3.4. *There exists a diffeomorphism of the Seifert fiber space X/π which realizes the group extension* (6) *over the base $\mathcal{O}$ to the orbifold M_i.*

In particular, the smooth manifold (respectively orbifold) M carries a Seifert fiber space structure over the orbifold $\mathcal{O} = Y/\Theta$.

3.3. **Collapsing of irreducible fake tori.** We show now that irreducible fake smooth tori do not collapse to aspherical Riemannian orbifolds.

Proposition 3.5. *Let $\mathcal{T}$ be an irreducible fake smooth torus. Then $\mathcal{T}$ does not admit a collapse with bounded curvature to a compact aspherical Riemannian orbifold $\mathcal{O} = Y/\Theta$.*

Proof. Assume that M has such a collapse. By Theorem 1.1, M admits the structure of a Seifert fiber space over $\mathcal{O}$ with infranil fiber, which realizes a group extension of the form

$$1 \to \mathbb{Z}^{n-k} \to \mathbb{Z}^n \to \Theta \to 1 \ .$$

Since Θ is abelian and it acts properly discontinuously with compact quotient on the contractible manifold Y, we infer that Θ is torsion-free abelian, isomorphic to $\mathbb{Z}^k$, $k < n$. Therefore, the above extension is isomorphic to the direct product

$$\mathbb{Z}^n = \mathbb{Z}^{n-k} \times \mathbb{Z}^k \, .$$

Note that $\mathcal{O} = Y/\Theta$ is a manifold and a homotopy torus. This implies that $\mathcal{O}$ is also homeomorphic to a torus T^k. The Seifert rigidity Theorem 2.8 implies that the Seifert fiber space $\mathcal{T}$ over $\mathcal{O}$ is affinely diffeomorphic to the product $T^{n-k} \times \mathcal{O}$. Therefore, $\mathcal{O}$ is a fake smooth torus. This contradicts the irreducibility of $\mathcal{T}$. □

We can give now the proof for Corollary 1.6:

Proof. Assume to the contrary that D-MinVol$(\mathcal{T}) = 0$. Since $\mathcal{T}$ is aspherical, this assumption implies (see the proof of Fukaya's Theorem [9, Theorem 0-1] or [7, Theorem 16.1]) that $\mathcal{T}$ admits a bounded curvature collapse to a compact aspherical Riemannian orbifold $\mathcal{O} = Y/\Theta$. This is not possible by the previous proposition. □

4. Seifert fiberings on infrasolv manifolds

Let S be a simply connected *solvable* Lie group.

Definition 4.1. A compact infra-S space S/Δ will be called an *infrasolv orbifold.* If S/Δ is a manifold, it is called an *infrasolv manifold.*

In this section we prove that up to diffeomorphism every infrasolv orbifold arises from a Seifert fiber construction with nil-geometry over a flat Riemannian orbifold. In addition we show that the class of manifolds which are diffeomorphic to infrasolv manifolds is closed with respect to the Seifert fiber construction with nil-geometry. In view of Theorem 1.1, this implies Theorem 1.3 in the introduction.

4.1. Flat orbifolds associated to virtually abelian groups.

Let $\bar{\Theta}$ be a finitely generated virtually abelian group of rank k. Then there exists a homomorphism $\bar{\Theta} \to \mathrm{Isom}(\mathbb{R}^k)$ such that $\bar{\Theta}$ acts properly discontinuously on $\mathbb{R}^k$. This homomorphism is unique up to conjugacy by affine transformations (that is, by elements of $\mathrm{Aff}(\mathbb{R}^k)$). (By Bieberbach's second and third theorems, see, for example, [24].) Let $\mathcal{O}_{\bar{\Theta}} = \mathbb{R}^k/\bar{\Theta}$ denote the flat orbifold associated to $\bar{\Theta}$.

4.2. Seifert fiberings over flat orbifolds.

The following implies Theorem 1.2 in the introduction:

Theorem 4.2. *Let $M = S/\pi$ be an infrasolv orbifold, and*

$$1 \longrightarrow \Delta_0 \longrightarrow \pi \longrightarrow \bar{\Theta} \longrightarrow 1 \tag{8}$$

an exact sequence of groups, where Δ_0 is a finitely generated torsion-free nilpotent group. Assume further that $\bar{\Theta}$ is a virtually abelian finitely generated group of rank k. Then M admits a Seifert fiber structure over the flat orbifold $\mathcal{O}_{\bar{\Theta}} = \mathbb{R}^k/\bar{\Theta}$, which realizes the group extension (8).

Proof. Since M is diffeomorphic to a standard π-orbifold M_π (see below), it is enough to show the statement for M_π. Therefore, the theorem is a consequence of Proposition 4.8 below. □

Corollary 4.3. *Every infrasolv orbifold has a Seifert fiber structure (with nil-geometry) over a flat orbifold.*

Proof. Indeed, we may consider the Fitting subgroup Δ_0 of π, which is (by definition) the maximal nilpotent normal subgroup of π. This gives rise to an exact sequence of the form (8). □

Let π be a torsion-free virtually polycyclic group. Then (by [2], but see also Section 4.3.4 below), there exists an infrasolv manifold M_π with fundamental group π.

Corollary 4.4. *Let X/π be the total space of a Seifert fiber space with nil-geometry over a flat orbifold $\mathcal{O} = \mathbb{R}^k/\Theta$. Then X/π is (diffeomorphic to) an infrasolv manifold.*

Proof. Since X/π is a Seifert fiber space over the base $\mathcal{O}$, there exists a homomorphism $\pi \to \Theta$, as in (8). Let M_π be an infrasolv manifold with fundamental group π, and $f : M_\pi \to \mathcal{O}$ the bundle map for the Seifert fiber structure on M_π which realizes the group extension representing the map $\pi \to \Theta$. By Theorem 2.8 (uniqueness of the Seifert construction), the infra-N bundles M_π and X/π are fiberwise affinely diffeomorphic over $\mathcal{O}$. In particular, M_π and X/π are diffeomorphic. □

4.3. **Flat affine manifolds.**

Let U be a unipotent real linear algebraic group. Then U is in particular a simply connected nilpotent Lie group. Conversely, every simply connected nilpotent Lie group admits the structure of a unipotent real linear algebraic group. (Cf. [19, Preliminaries, §2] for basic facts on linear algebraic groups.)

4.3.1. *Flat manifolds modeled on U.* We consider manifolds which carry locally the *affine* geometry of U. This generalizes the notion of infra-U manifold, which is an essentially metric concept.

Let $\mathrm{Aff}(U) = \mathrm{Aff}(U, \nabla^{\mathrm{can}})$ be the affine group of U.

Definition 4.5. Let $\Delta \leq \mathrm{Aff}(U)$ be a discrete subgroup which acts properly on U with compact quotient. Then the quotient

$$U/\Delta$$

is called a *flat affine U-orbifold.* If Δ acts freely on U then U/Δ is a *flat affine U-manifold.*

Remark. A long standing conjecture ("Auslander's conjecture") asserts that Δ must be a virtually polycyclic group.

4.3.2. *Unipotent shadow and algebraic hull.* Now let Δ be a virtually polycyclic group without a finite (non-trivial) normal subgroup. By a construction, originally due to Mostow and Auslander (see [19, Chapter IV, Proposition 4.4] and [1, Appendix A.1], for reference), we may associate to Δ in a unique way a unipotent real linear algebraic group U_Δ, which is called the *unipotent shadow* of Δ. The unipotent shadow U_Δ has the following characterizing additional properties:

(1) $\dim U_\Delta = \operatorname{rank} \Delta$. (Recall that the rank of Δ coincides with the virtual cohomological dimension of Δ.)
(2) There exists a $\mathbb{Q}$-defined real linear algebraic group $H = H_\Delta$ which has $\Delta \leq H(\mathbb{Q})$ as a Zariski-dense subgroup.
(3) There is a splitting of algebraic groups $H = U_\Delta \rtimes T$, such that T is a subgroup of semi-simple elements which acts faithfully on U_Δ.

The group H is called the *algebraic hull of* Δ.

4.3.3. *Standard Δ-manifolds.* Since T acts faithfully on U_Δ, we may realize the hull $H = H_\Delta$ as a subgroup of $\operatorname{Aff}(U_\Delta)$ in a natural way. In particular, in this way, Δ acts faithfully on U_Δ by affine transformations. Then we have:

Theorem 4.6 (cf. [1, Theorem 1.2])**.** *The affine action of Δ on U_Δ is properly discontinuous, and it is uniquely defined up to conjugacy with an element of the affine group* $\operatorname{Aff}(U)$*.*

The quotient space

$$M_\Delta = U_\Delta \big/ \Delta$$

is called a *standard Δ-orbifold.* If M_Δ is a manifold it is called a *standard Δ-manifold.* Remarkably, in standard Δ-manifolds all homotopy equivalences of maps are induced by affine automorphisms of U_Δ, see [1].

Remark. If $\bar{\Delta}$ is an arbitrary virtually polycyclic group, dividing by its maximal finite normal subgroup yields a unique homomorphism $\bar{\Delta} \to \Delta$ onto the fundamental group of a standard Δ-orbifold. We also call $M_{\bar{\Delta}} = M_\Delta$ the standard $\bar{\Delta}$-orbifold associated to $\bar{\Delta}$.

4.3.4. *Smooth rigidity of infrasolv manifolds.* As remarked in [1], every standard Δ-manifold admits a (in general, non-unique) infrasolv manifold structure modeled on some solvable Lie group. In particular, every torsion-free group Δ as above appears as the fundamental group of an infrasolv manifold. Moreover, standard Δ-manifolds may serve as *unique* smooth models for infrasolv manifolds:

Theorem 4.7 (cf. [1, Theorem 1.4])**.** *Let $M = S/\Delta$ be a (compact) infrasolv manifold, where S is a simply connected solvable Lie group and $\Delta \leq \mathrm{Aff}(S)$ a discrete group of isometries. Then M is smoothly diffeomorphic to the standard Δ-manifold M_Δ.*

The corresponding statements for infrasolv orbifolds and standard Δ-orbifolds hold as well, with the same proof as in [1].

4.4. **Seifert fiberings on standard-manifolds.** Let M_π be a standard π-orbifold, where π is a virtually polycyclic group. Consider an exact sequence of groups

$$1 \longrightarrow \Delta_0 \longrightarrow \pi \longrightarrow \bar{\Theta} \longrightarrow 1 \,, \tag{9}$$

where Δ_0 is a finitely generated torsion-free nilpotent group.

Proposition 4.8. *There exists a Seifert fiber structure for M_π, which realizes the group extension* (9) *over the standard $\bar{\Theta}$-orbifold $M_{\bar{\Theta}}$.*

Proof. Let $U = U_\pi$ be the unipotent shadow of π. Recall that for the standard affine embedding $\pi \leq \mathrm{Aff}(U)$ the intersection $U \cap \pi$ is the maximal nilpotent normal subgroup $\mathrm{Fitt}(\pi)$ of π (see [1]). Since Δ_0 is nilpotent, we therefore have $\Delta_0 \leq U$. Let U_0 be the Zariski-closure of Δ_0 in U. Then Δ_0 is a finite index subgroup of $\Delta_0' = U_0 \cap \pi$. Moreover, U_0 is normalized by π, since Δ_0 is a normal subgroup of π. Since π is Zariski-dense in $H = UT$ it follows, in particular, that U_0 is normal in H and hence also in U. Let $U_1 = U/U_0$ be the quotient unipotent $\mathbb{Q}$-defined real linear algebraic group.

We consider now the induced homomorphism $H \to \mathrm{Aff}(U_1)$. Let Θ denote the image of π in $\mathrm{Aff}(U_1)$. It clearly is a Zariski dense subgroup of the image H_1 of H in $\mathrm{Aff}(U_1)$. Since U_1 is the unipotent radical of H_1, we have the relation $\dim U - \dim U_0 = \dim U_1 \leq \operatorname{rank} \Theta$ (see [19, Chapter IV, Lemma 4.36]). Since Δ_0 is contained in the kernel of $\pi \to \Theta$, we have $\operatorname{rank} \Theta \leq \operatorname{rank} \pi - \operatorname{rank} \Delta_0 = \dim U - \dim U_0$. We conclude that $\operatorname{rank} \Theta = \dim U_1$. It is now evident that the image H_1 of H in $\mathrm{Aff}(U_1)$ satisfies the axioms for an algebraic hull of Θ. Therefore $M_\Theta = U_1/\Theta$ is a standard Θ-manifold. The data $(X, N, \pi) = (U, U_0, \pi)$ define a Seifert bundle map

$$\sigma : M_\pi \to M_\Theta$$

and the Seifert construction (U, U_0, π) realizes the group extension (9) over $M_{\bar{\Theta}} = M_\Theta$. □

Corollary 4.9. *The class of smooth orbifolds represented by standard Δ-orbifolds is closed with respect to the Seifert fiber construction with nil-geometry.*

Proof. Indeed, let $\sigma : M \to M_{\bar{\Theta}}$ be a Seifert fiber space over a standard orbifold $M_{\bar{\Theta}}$, which realizes a group extension of the form (9). By the previous proposition, the standard-π orbifold M_π also supports a Seifert construction with nil-geometry over $M_{\bar{\Theta}}$, which realizes (9). By the Seifert rigidity theorem (Theorem 2.8), M and M_π are diffeomorphic. □

5. APPENDIX

In the following sections we gather several important notions and results from equivariant Gromov-Hausdorff convergence, developed by Fukaya and Yamaguchi, and the work of Cheeger, Fukaya, and Gromov on collapsed Riemannian manifolds with bounded sectional curvature. General references for these topics are [7] and [15].

Appendix A. (Equivariant) Gromov-Hausdorff Convergence

Let us first recall the definition of the classical Hausdorff distance: For subsets A and B of a metric space X, the *Hausdorff distance* between A and B in X, $d_H^X(A, B)$, is defined as the infimum of all positive real numbers ε such that the open ε-neighbourhood of A in X contains B and vice versa. For any metric space X, the Hausdorff distance d_H^X then defines a metric on the set of all closed and bounded nonempty subsets of X.

Around 1980, Gromov introduced an abstract version of the Hausdorff distance as follows: The *Gromov-Hausdorff distance* of two compact metric spaces X and Y, $d_{GH}(X, Y)$, is defined as the infimum of all numbers $d_H^Z(f(X), g(Y))$, where Z ranges over all metric spaces in which X and Y can be imbedded isometrically, and f and g over all isometric embeddings $X \to Z$ and $Y \to Z$. Alternatively, the Gromov-Hausdorff distance of X and Y can also be defined as the infimum of all Hausdorff distances $d_H^Z(X, Y)$, where Z is the disjoint union of X and Y and where the infimum is now taken over all metrics on Z which extend the metrics on X and Y.

The Gromov-Hausdorff distance d_{GH} defines a metric on the collection $\mathcal{C}$ of all compact metric spaces, considered up to isometry, and therefore gives rise to the notion of *Gromov-Hausdorff convergence* of sequences of compact metric spaces. Moreover, with respect to d_{GH}, the space $\mathcal{C}$ is complete.

It is sometimes also convenient to work with so-called (Gromov-) Hausdorff approximations: If X and Y are compact metric spaces and ε is real and positive, a (not necessarily continuous) map $\phi : X \to Y$ is called an *$\varepsilon-$Hausdorff approximation from X to Y* if $|\, d_Y(\phi(x), \phi(x')) - d_X(x, x')\,| \leq \varepsilon$ for all $x, x' \in X$ and if the $\varepsilon-$neighbourhood of $\phi(X)$ in Y is equal to Y.

Notice that if $d_{GH}(X, Y)$ is less than ε, then there exist $3\varepsilon-$Hausdorff approximations from X to Y and Y to X. Conversely, if there exist $\varepsilon-$Hausdorff approximations between X and Y, then $d_{GH}(X, Y) \leq 3\varepsilon$. In particular, Gromov-Hausdorff convergence can therefore also be defined via Hausdorff approximations.

There is also a useful notion of Gromov-Hausdorff convergence for noncompact metric spaces, at least for *length spaces* in which, by definition, the distance between points is given by the infimum of the lengths of all curves joining the points:

A sequence $(X_n, x_n)_{n\in\mathbb{N}}$ of locally compact length spaces with given basepoints $x_n \in X_n$ is said to converge to a pointed metric space (Y, y) in the pointed Gromov-Hausdorff sense if, for all $r > 0$, the closed r-balls $B_r(x_n)$ Gromov-Hausdorff converge in the usual sense to the closed r-ball $B_r(y)$ in Y.

Under this so-called *pointed Gromov-Hausdorff convergence* or *Gromov-Hausdorff convergence with basepoint*, the collection $\mathcal{M}$ of all locally compact complete pointed length spaces, considered up to isometries that preserve basepoints, is complete as well. Moreover, the same is true for ordinary Gromov-Hausdorff convergence in the subspaces $\mathcal{L}(D)$ of $\mathcal{C}$ which are made up of the isometry classes of compact length spaces subject to a given uniform upper bound D for the diameter.

In the context of Riemannian manifolds with lower curvature bounds Gromov obtained the following fundamental result:

Theorem A.1 (Gromov's Precompactness Theorem) *For any natural number m and any real numbers κ and D, the class of closed Riemannian m-manifolds with Ricci curvature* $\mathrm{Ric} \geq (m-1)\,\kappa$ *and diameter* $\mathrm{diam} \leq D$ *is precompact in $\mathcal{L}(D)$ with respect to Gromov-Hausdorff convergence, and the class of pointed complete Riemannian m-manifolds with $Ric \geq (m-1)\,\kappa$ is precompact in $\mathcal{M}$ with respect to the pointed Gromov-Hausdorff topology.*

If one replaces the Ricci curvature bounds in Theorem A.1 by sectional curvature ones, one may actually replace $\mathcal{L}(D)$ and $\mathcal{M}$ by appropiate classes of Alexandrov spaces, but we will not need this here.

Instead, let us now turn to notions of Gromov-Hausdorff convergence for metric spaces which are equipped with isometric group actions.

Let $\mathcal{M}$ be as above, and let $\mathcal{M}^{Groups}$ consist of all triples (X, Γ, p) with $(X, p) \in \mathcal{M}$ such that Γ is a closed subgroup of the isometry group of X, and let in this case for real positive r be $\Gamma(r)$ the set of all $\gamma \in \Gamma$ for which $d(\gamma p, p) < r$.

Definition A.2 *An* ε-equivariant Hausdorff approximation with basepoint p *from* $(X, \Gamma, p) \in \mathcal{M}^G$ *to* $(Y, \Lambda, q) \in \mathcal{M}^{Groups}$ *is given by a triple of maps* (f, ϕ, ψ) *with*

$$f : B_p(1/\varepsilon, X) \to Y, \quad \phi : \Gamma(1/\varepsilon) \to \Lambda(1/\varepsilon) \quad \text{and} \quad \psi : \Lambda(1/\varepsilon) \to \Gamma(1/\varepsilon)$$

that satisfies the following conditions:

1. $f(p) = q$;
2. *the* ε*-neighbourhood of* $f(B_p(1/\varepsilon, X))$ *contains* $B_q(1/\varepsilon, Y)$;
3. *for all* $x_1, x_2 \in B_p(1/\varepsilon, X)$ *one has*
$$|\, d(f(x_1), f(x_2)) - d(x_1, x_2)\, | < \varepsilon \,;$$
4. *if* $\gamma \in \Gamma(1/\varepsilon)$, $x \in B_p(1/\varepsilon, X)$, *and* $\gamma\, x \in B_p(1/\varepsilon, X)$, *then*
$$d\,(f(\gamma\, x), \phi(\gamma)\,(f(x))) < \varepsilon \,;$$
5. *if* $\mu \in \Lambda(1/\varepsilon)$, $x \in (B_p(1/\varepsilon, X)$, *and* $\psi(\mu)\,(x) \in (B_p(1/\varepsilon, X)$, *then*
$$d\,(f(\psi(\mu)\,(x)), \mu\,(f(x)) < \varepsilon \,.$$

Notice that it is neither required that the maps occuring here be globally defined, nor that they be continuous, nor, as regards ϕ and ψ, be homomorphisms.

Definition A.3 *Let* $(X, \Gamma, p), (Y, \Lambda, q) \in \mathcal{M}^{Groups}$. *The* equivariant Gromov-Hausdorff distance with basepoint $d^e_{GH}((X, \Gamma, p), (Y, \Lambda, q))$ *is defined as the infimum of all positive real numbers* ε *such that there exist* ε*-equivariant Hausdorff approximations with basepoint from* (X, Γ, p) *to* (Y, Λ, q) *and vice versa.*

If $(X_n, \Gamma_n, p_n), (Y, \Lambda, q) \in \mathcal{M}^{Groups}$*, the equation*

$$\lim_{n\to\infty} \quad (X_n, \Gamma_n, p_n) = (Y, \Lambda, q)$$

means that

$$\lim_{n\to\infty} \quad d^e_{GH}((X_n, \Gamma_n, p_n), (Y, \Lambda, q)) = 0\,.$$

If the groups that occur in the above definition are isomorphic to a fixed group G, we will also speak of *G-equivariant Gromov-Hausdorff distance (with basepoint).*

In the case where all groups are trivial, we will write (X, p) instead of (X, id, p) etc. and d_{GH} instead of d^e_{GH}, and equivariant Gromov-Hausdorff convergence with basepoint then reduces to ordinary Gromov-Hausdorff convergence with basepoint.

Let us also note that in the situtation where all spaces in question are compact and have uniformly bounded diameters, Gromov-Hausdorff convergence and Gromov-Hausdorff convergence with base point are essentially equivalent concepts: The latter clearly implies the former, and up to passing to a subsequence the converse also holds.

Let us now gather several important facts and results concerning equivariant Gromov-Hausdorff convergence.

Proposition A.4 ([13]) *If* $(X_n, \Gamma_n, p_n) \in \mathcal{M}^{Groups}$ *and* $(Y, q) \in \mathcal{M}$ *satisfy with respect to the Gromov-Hausdorff distance with basepoint*

$$\lim_{n\to\infty} \quad (X_n, p_n) = (Y, q)\,,$$

then there exist Λ *and a subsequence* $\{n_k\}_{k\in\mathbb{N}}$ *such that* $(Y, \Lambda, q) \in \mathcal{M}^{Groups}$ *and such that*

$$\lim_{k\to\infty} \quad (X_{n_k}, \Gamma_{n_k}, p_{n_k}) = (Y, \Lambda, q)\,.$$

Notice that even if all Γ_n and X_n above are the same, in general the limit group Λ can be a different group. However, for compact groups and spaces we have the following stability result.

Proposition A.5 ([7, Theorem 6.9]) *Let* G *be a group and* $\mathcal{M}(G)$ *denote the set of all isometry classes of compact metric spaces equipped with an isometric* G*-action. If* G *is compact, then for any natural number* m *and any real numbers* κ *and* D*, the class of closed* m*-dimensional Riemannian* G*-manifolds with Ricci curvature* $\mathrm{Ric} \geq (m-1)\,\kappa$ *and* *diameter* diam $\leq D$ *is precompact in* $\mathcal{M}(G)$ *with respect to the* G-*equivariant Gromov-Hausdorff topology.*

The following results establish several relations between equivariant Gromov-Hausdorff convergence and convergence of subgroups and quotient spaces.

Proposition A.6 ([12]) *If* $(X_n, \Gamma_n, p_n), (Y, \Lambda, q) \in \mathcal{M}^{Groups}$ *and*

$$\lim_{n\to\infty} \quad (X_n, \Gamma_n, p_n) = (Y, \Lambda, q)\,,$$

then

$$\lim_{n\to\infty} \quad (X_n/\Gamma_n, \bar{p}_n) = (Y/\Lambda, \bar{q})\,,$$

where $\bar{p}_n$ *and* $\bar{q}$ *denote the images of* p_n *and* q*, respectively, under the canonical orbit space projections* $X_n \to X_n/\Gamma_n$ *and* $Y \to Y/\Lambda$*.*

Proposition A.7 ([12]) *Let* X_n *and* Y *be complete and simply connected Riemannian manifolds and* Γ_n *resp.* Λ *be closed subgroups of the isometry groups of* X_n *resp.* Y *which all act effectively and properly discontinuously. Suppose, moreover, that the sectional curvatures of* X_n *and* Y *are uniformly bounded in absolute value. Then*

$$\lim_{n\to\infty} \quad (X_n/\Gamma_n, \bar{p}_n) = (Y/\Lambda, \bar{q})$$

implies

$$\lim_{n\to\infty} \quad (X_n, \Gamma_n, p_n) = (Y, \Lambda, q)\,.$$

Proposition A.8 ([13]) *Let* $(X_n, \Gamma_n, p_n), (Y, \Lambda, q) \in \mathcal{M}^{Groups}$ *satisfy*

$$\lim_{n\to\infty} \quad (X_n, \Gamma_n, p_n) = (Y, \Lambda, q)$$

and let Λ' *be a normal subgroup of* Λ *such that the following conditions hold:*

1. Λ/Λ' *is discrete and finitely presented;*

2. Y/Λ *is compact;*

3. there exists $r > 0$ *such that* Λ' *is generated by* $\Lambda'(r)$ *and such that the homomorphism* $\pi_1(B_q(r, Y), q) \to \pi_1(Y, q)$ *is surjective;*

4. for all n*,* X_n *is simply connected and* Γ_n *acts freely and properly discontinuously on* X_n*.*

Then there exists a sequence of normal subgroups Γ'_n *of* Γ_n *satisfying*

$$(a) \quad \lim_{n\to\infty} \quad (X_n, \Gamma'_n, p_n) = (Y, \Lambda', q)\,;$$

(b) Γ_n/Γ'_n *is isomorphic to* Λ/Λ' *for all sufficiently large* n.

APPENDIX B. COLLAPSING OF MANIFOLDS UNDER BOTH-SIDED BOUNDS ON SECTIONAL CURVATURE

We will now briefly describe the collapsing of Riemannian manifolds with both-sided bounds on sectional curvature, where a detailed structure theory has been initiated and developed in the works of Cheeger, Fukaya, Gromov, and others.

Let $M = (M^m, g)$ be a Riemannian manifold of dimension m and let $FM = F(M^m)$ denote its bundle of orthonormal frames. When fixing a bi-invariant metric on $O(m)$, the Levi-Civita connection of g gives rise to a canonical metric on FM, so that the projection $FM \to M$ becomes a Riemannian submersion and so that $O(m)$ acts on FM by isometries. Another fibration structure on FM is called $O(m)$ *invariant* if the $O(m)$ action on FM preserves both its fibres and its structure group.

A *pure N-structure* on M^m is defined by an $O(m)$ invariant fibration, $\tilde{\eta} : FM \to B$, with fibre a nilmanifold isomorphic to $(N/\Gamma, \nabla^{\mathrm{can}})$ and structural group contained in the group of affine automorphisms of the fibre, where B is a smooth manifold, N is a simply connected nilpotent group and ∇^{can} the canonical connection on N for which all left invariant vector fields are parallel. A pure N-structure on M induces, by $O(m)$-invariance, a partition of M into "orbits" of this structure, and is then said to have *positive rank* if all these orbits have positive dimension. A pure N-structure $\tilde{\eta} : FM \to B$ over a Riemannian manifold (M, g) gives rise to a sheaf on FM whose local sections restrict to local right invariant vector fields on the fibres of $\tilde{\eta}$, and if the local sections of this sheaf are local Killing fields for the metric g, then g is said to be *invariant* for the N-structure (and $\tilde{\eta}$ is then also sometimes referred to as *pure nilpotent Killing structure for* g). Cheeger, Fukaya and Gromov showed:

Theorem B.1 ([4]) *Let for* $m \geq 2$ *and* $D > 0$ $\mathfrak{M}(m, D)$ *denote the class of all m-dimensional compact connected Riemannian manifolds* (M, g) *with sectional curvature* $|K_g| \leq 1$ *and diameter* $diam(g) \leq D$.

Then, given any $\varepsilon > 0$*, there exists a positive number* $v = v(m, D, \varepsilon) > 0$ *such that if* $(M, g) \in \mathfrak{M}(m, D)$ *satisfies* $vol(g) < v$*, then* M^m *admits a pure N-structure* $\tilde{\eta} : FM \to B$ *of positive rank so that*

(a) there is a smooth metric g_ε *on* M *which is invariant for the N-structure* $\tilde{\eta}$ *and for which all fibres of* $\tilde{\eta}$ *have diameter less than* ε*, satisfying*

$$e^{-\varepsilon} g < g_\varepsilon < e^{\varepsilon} g \ , \qquad |\, \nabla_g - \nabla_{g_\varepsilon} \,| < \ \varepsilon \ , \qquad |\, \nabla^l_{g_\varepsilon} R_{g_\varepsilon} \,| < \ C(m, l, \varepsilon) \ ;$$

(b) there exist constants $i = i(m, \varepsilon) > 0$ *and* $C = C(m, \varepsilon)$ *such that, when equipped with the metric induced by* g_ε*, the injectivity radius of* B *is* $\geq i$ *and the second fundamental form of all fibres of* $\tilde{\eta}$ *is bounded by* C*.*

The $O(m)$ invariance of a pure N-structure $\tilde{\eta} : FM \to B$ implies that the $O(m)$ action on FM descends to an $O(m)$ action on B and that the fibration on FM descends to a possibly singular fibration on M, $\eta : M^m \to B/O(m)$, such that the following diagram commutes.

$$\begin{tikzcd}
F(M^m) \arrow[r, "\tilde{\eta}"] \arrow[d, "\pi"] & B \arrow[d, "\tilde{\pi}"] \\
M^m \arrow[r, "\eta"] & B/O(m)
\end{tikzcd}$$

Theorem B.1 describes the structure of a general collapse with bounded curvature, where $B/O(m)$ is in general not a manifold. However, in the case where one has a bounded curvature collapse to a manifold limit space, there is also Fukaya's following equivariant fibre bundle version of the above result:

Theorem B.2 ([7, 8, 10]) *Let* $M_i \in \mathfrak{M}(m, D)$*,* $N \in \mathfrak{M}(n, D)$ *and let* G *be a compact Lie group which acts on* M_i *and* N *by isometries so that the* M_n *converge for* $i \to \infty$ *to* N *in the* G*-equivariant Gromov-Hausdorff distance. Then, for i sufficiently large, there exist mappings* $\pi_i : M_i \to N$ *which satisfisfy*

(a) $\pi_i : M_i \to N$ *is a fibre bundle and* π_i *is* G*-equivariant;*

(b) each fiber $\pi_i^{-1}(p)$ *carries a flat connection which depends smoothly on* $p \in N$*, such that* $\pi_i^{-1}(p)$ *is affinely diffeomorphic to an infranil manifold* N_i/Δ_i *with its canonical flat connection* ∇^{can}*;*

(c) the affine structures on the fibres of $\pi_i : M_i \to N$ *are* G*-invariant;*

(d) the structure group of π_i *is contained in the Lie group*

$$\frac{\mathbb{C}(N_i)}{\mathbb{C}(N_i) \cap \Delta_i} \rtimes \mathrm{Aut}(\Delta_i)\,,$$

where $\mathbb{C}(N_i)$ *denotes the center of* N_i*;*

(e) π_i *is an almost Riemannian submersion in the sense that for any* $p \in N$ *and any tangent vector* $v \in T_pM_i$ *orthogonal to the fibre* $\pi_i^{-1}(p)$*, one has an estimate*

$$e^{-o(i)} \quad < \quad ||d\pi_i(v)|| \;/\; ||v|| \quad < \quad e^{o(i)}\,,$$

where $o(i)$ *satisfies* $lim_{i\to\infty} o(i) = 0$*.*

References

[1] O. Baues, *Infra-solvmanifolds and rigidity of subgroups in solvable linear algebraic groups*, Topology, **43** (2004), no. 4, 903-924.

[2] L. Auslander, F.E.A. Johnson, *On a conjecture of C.T.C. Wall*, J. London Math. Soc. **14** (1972), no. 2, 331-332.

[3] W. Browder, *On the action of* $\Theta^n\ (\partial\pi)$, Differential and Combinatorial Topology (A Symposium in Honor of Marston Morse), 23-36, Princeton Univ. Press, Princeton, N.J., 1965.

[4] J. Cheeger, K. Fukaya, M. Gromov, *Nilpotent structures and invariant metrics on collapsed manifolds*, J. Amer. Math. Soc. **5** (1992), no. 2, 327-372.

[5] J. Cheeger, X. Rong, *Existence of polarized F-structures on collapsed manifolds with bounded curvature and diameter*, Geom. Funct. Anal. **6** (1996), no. 3, 411-429.

[6] F. T. Farrell, W. C. Hsiang, *Topological characterization of flat and almost flat Riemannian manifolds* M^n $(n \neq 3, 4)$, *Amer. J. Math.* **105** (1983), 641-672.

[7] K. Fukaya, *Hausdorff convergence of Riemannian manifolds and its applications*, Recent Topics in Differential and Analytic Geometry, Advanced Studies in Pure Mathematics **18**-I, 143-238, 1990.

[8] K. Fukaya, *Collapsing Riemannian manifolds to ones with lower dimension. II.* J. Math. Soc. Japan **41** (1989), no. 2, 333-356.

[9] K. Fukaya, *A compactness theorem of a set of aspherical Riemannian orbifolds*, A fête of topology, 391-413, Academic Press, Boston, MA, 1988.

[10] K. Fukaya, *A boundary of the set of the Riemannian manifolds with bounded curvatures and diameters*, J. Differential Geom. **28** (1988), no. 1, 1-21.

[11] K. Fukaya, *Collapsing Riemannian manifolds to ones of lower dimensions*, J. Differential Geom. **25** (1987), no. 1, 139-156.

[12] K. Fukaya, *Theory of convergence for Riemannian orbifolds*, Japan J. Math. **12** (1986), 121-160.

[13] K. Fukaya and T. Yamaguchi. *The fundamental groups of almost nonnegatively curved manifolds*, Ann. of Math. (2) **136** (1992), 253-333.

[14] M. Gromov, *Almost flat manifolds*, J. Differential Geom. **13** (1978), 231-241.

[15] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces.* Progress in Mathematics, vol. 152, Birkhäuser, Boston, 1999.

[16] Y. Kamishima, K.B. Lee and F. Raymond, *The Seifert construction and its applications to infranil manifolds*, Quart. J. Math., Oxford (2), **34** (1983), 433-452.

[17] K.B. Lee and F. Raymond, *Seifert manifolds*, Handbook of geometric topology, 635-705, North-Holland, Amsterdam, 2002.

[18] K.B. Lee and F. Raymond, *Seifert fiberings.* Mathematical Surveys and Monographs, 166. American Mathematical Society, Providence, RI, 2010.

[19] M.S. Raghunathan, *Discrete subgroups of Lie groups*, Ergebnisse Math. vol. 68, Springer Verlag, Berlin, New York, 1972.

[20] X. Rong, *The existence of polarized F-structures on volume collapsed 4-manifolds*, Geom. Funct. Anal. **3** (1993), no. 5, 474-501.

[21] E.A Ruh, *Almost flat manifolds*, J. Differential Geom. **17** (1982), 1-14.

[22] W. Tuschmann, *Collapsing, solvmanifolds and infrahomogeneous spaces*, Differential Geom. Appl. 7 (1997), no. 3, 251-264.

[23] C.T.C. Wall, *Surgery on compact manifolds*, Academic Press, 1970.

[24] J. A. Wolf, *Spaces of constant curvature*, new edition, AMS 2010.

Oliver Baues and Wilderich Tuschmann, Fakultät für Mathematik, Karlsruher Institut für Technologie (KIT), D-76128 Karlsruhe, Germany

E-mail address: `oliver.baues@kit.edu`

E-mail address: `tuschmann@kit.edu`